

- Chakradhari, S., Perkons, I., Mišina, I., **Sipeniece, E.**, Radziejewska-Kubzdela, E., Grygier, A., Rudzińska, M., Patel, K. S., Radzimirska-Graczyk, M., & Górnaś, P. (2020). Profiling of the bioactive components of safflower seeds and seed oil: Cultivated (*Carthamus tinctorius* L.) vs. wild (*Carthamus oxyacantha* M. Bieb.). *European Food Research and Technology*, 246(3), 449–459. <https://doi.org/10.1007/s00217-019-03414-w>
- Fatty Acids, Tocopherols, Tocotrienols, Phytosterols, Carotenoids, and Squalene in Seed Oils of Hyptis suaveolens, Leonotis nepetifolia, and Ocimum sanctum—Sahu—2020—European Journal of Lipid Science and Technology—Wiley Online Library.* (n.d.). Retrieved 6 June 2024, from <https://onlinelibrary.wiley.com/doi/full/10.1002/ejlt.202000053>
- Górnaś, P., Czubinski, J., Rudzińska, M., Grygier, A., Ying, Q., Chakradhari, S., Sahu, P. K., Mišina, I., **Urvaka, E.**, & Patel, K. S. (2019). Selected Uncommon Legumes as a Source of Essential Fatty Acids, Tocopherols, Tocotrienols, Sterols, Carotenoids, and Squalene. *Plant Foods for Human Nutrition*, 74(1), 91–98. <https://doi.org/10.1007/s11130-018-0706-x>
- Górnaś, P., Rudzińska, M., Grygier, A., Ying, Q., Mišina, I., **Urvaka, E.**, & Rungis, D. (2019). Sustainable valorization of oak acorns as a potential source of oil rich in bioactive compounds. *Process Safety and Environmental Protection*, 128, 244–250. <https://doi.org/10.1016/j.psep.2019.06.003>
- Mišina, I., **Sipeniece, E.**, Grygier, A., Qian, Y., Rudzińska, M., Kaufmane, E., Segliņa, D., Siger, A., & Górnaś, P. (2021). Profiling of the lipophilic components of seed oils recovered from twelve Japanese quince (*Chaenomeles japonica*) genotypes. *Natural Product Research*, 36(1), 474–478. <https://doi.org/10.1080/14786419.2020.1782407>
- Mišina, I., **Sipeniece, E.**, Rudzińska, M., Grygier, A., Radzimirska-Graczyk, M., Kaufmane, E., Segliņa, D., Lācis, G., & Górnaś, P. (2020). Associations between Oil Yield and Profile of Fatty Acids, Sterols, Squalene, Carotenoids, and Tocopherols in Seed Oil of Selected Japanese Quince Genotypes during Fruit Development. *European Journal of Lipid Science and Technology*, 122(4), 1900386. <https://doi.org/10.1002/ejlt.201900386>
- Sahu, P. K., **Sipeniece, E.**, Radziejewska-Kubzdela, E., Grygier, A., Patel, K. S., Rudzińska, M., Radzimirska-Graczyk, M., & Górnaś, P. (2021). Profiling of the lipophilic components of seed oil obtained from two medicinal plants *Corchorus capsularis* L. and *Hibiscus sabdariffa* L. *Natural Product Research*, 36(1), 450–454. <https://doi.org/10.1080/14786419.2020.1777119>
- Sipeniece, E.**, Mišina, I., Grygier, A., Qian, Y., Rudzińska, M., Kaufmane, E., Segliņa, D., Siger, A., & Górnaś, P. (2021). Impact of the harvest year of three cultivars of Japanese quince (*Chaenomeles japonica*) on the oil content and its composition. *Scientia Horticulturae*, 275, 109683. <https://doi.org/10.1016/j.scienta.2020.109683>
- Sipeniece, E.**, Mišina, I., Qian, Y., Grygier, A., Sobieszcańska, N., Sahu, P. K., Rudzińska, M., Patel, K. S., & Górnaś, P. (2021). Fatty Acid Profile and Squalene, Tocopherol, Carotenoid, Sterol Content of Seven Selected Consumed Legumes. *Plant Foods for Human Nutrition*, 76(1), 53–59. <https://doi.org/10.1007/s11130-020-00875-3>